arXiv:1103.4722v1 [math.OC] 24 Mar 2011

An Approach to the Minimization of the Mumford–Shah Functional using Γ-convergence and Topological Asymptotic Expansion

Markus Grasmair,[1] Monika Muszkieta,[2] and Otmar Scherzer[1,3]

[1]Computational Science Center
University of Vienna
Nordbergstr. 15, 1090 Wien, Austria

[2]Institute of Mathematics and Computer Science
Wroclaw University of Technology
ul. Wybrzeze Wyspianskiego 27, 50-370 Wroclaw, Poland

[3]RICAM
Austrian Academy of Sciences
Altenbergerstr. 69, 4040 Linz, Austria

markus.grasmair@univie.ac.at
monika.muszkieta@pwr.wroc.pl
otmar.scherzer@univie.ac.at

March 21, 2011

Abstract

In this paper, we present a method for the numerical minimization of the Mumford–Shah functional that is based on the idea of topological asymptotic expansions. The basic idea is to cover the expected edge set with balls of radius $\varepsilon > 0$ and use the number of balls, multiplied with 2ε, as an estimate for the length of the edge set. We introduce a functional based on this idea and prove that it converges in the sense of Γ-limits to the Mumford–Shah functional. Moreover, we show that ideas from topological asymptotic analysis can be used for determining where to position the balls covering the edge set. The results of the proposed method are presented by means of two numerical examples and compared with the results of the classical approximation due to Ambrosio and Tortorelli.

1 Introduction

Let Ω be a Lipschitz bounded open domain in $\mathbb{R}^2$. We assume that a possibly noisy image on Ω is given, represented by a real-valued, bounded function f on Ω, whose values $f(x)$, $x \in \Omega$, correspond to the intensity of f at the pixel x. In order to segment, and denoise at the same time, the image f, Mumford and Shah [36] introduced a variational model, which is based on the assumptions that the different "objects" in the image give rise to homogeneous regions that are separated by the objects' projected silhouettes. Moreover, these silhouettes in general correspond to discontinuities in the image f. By this reasoning, they proposed to minimize the functional

$$F(u,K) = \frac{1}{2}\int_\Omega (u-f)^2\,dx + \frac{\alpha}{2}\int_{\Omega\setminus K} |\nabla u|^2\,dx + \mathcal{H}^1(K)\,, \tag{1.1}$$

taking as variables the function $u \in L^2(\Omega)$, the denoised image, and the compact set $K \subset \Omega$, the set of edges or silhouettes. Here, $\mathcal{H}^1(K)$ denotes the one-dimensional Hausdorff measure of the set K; in the case K is a regular (rectifiable) one-dimensional set, this is precisely its length. The parameters α and β that appear in (1.1) are positive constants determining the weight that is put on the regularity of the denoised image u and the length of the edge set K.

In order to show the existence of minimizers of the Mumford–Shah functional, a weak formulation depending only on one variable has been introduced by De Giorgi et al. [26]. In their model, the set K is replaced by the jump set S_u of the function $u \in SBV(\Omega)$, the space of special function of bounded variation on Ω. Still, this reformulation of the functional provides no method for the actual numerical computation of minimizers. Thus, various approximations of the functional F have been proposed, most of them based on the concept of Γ-convergence. Ambrosio and Tortorelli [3] proposed a variational model in which they replace the set S_u by a continuous function v with values close to 0 near S_u, and values close to 1 away from S_u. For other approximations of the Mumford–Shah functional in the sense of Γ-convergence, we refer to Braides et al. [13], where the authors propose approximations by a family of non-local functionals. Approximations by finite-difference schemes, inspired by the original discrete model of Blake and Zisserman [10], have been considered by Chambolle [16, 17], and by finite-elements schemes by Chambolle and Dal Maso [18] or recently by Aubert et al. [6]. In the work by Koepfler et al. [32] and Dal Maso et al. [25], region growing and merging methods have been proposed. For detailed analysis of the Mumford–Shah model, we refer reader to the book by Morel and Solimini [35]. We also refer to the books [7, 39, 40], where some of the above mentioned results are shortly discussed.

In the following, we will show how the problem of minimizing the Mumford–Shah functional F can be approached by means of an asymptotic expansion that is based on the idea of topological asymptotic analysis. In its original formulation (see, e.g., Sokołowski et al. [41], Garreau et al. [30], Feijóo et al. [29]), this theory investigates a variation of some objective functional depending on subsets of $\mathbb{R}^n$ with respect to the subtraction of a small ball from the considered domain. This variation is a scalar function, called the topological gradient or the topological derivative, and its largest negative values indicate positions, where it is good to remove a small ball. In [5], Amstutz proposed to modify the definition of topological gradient and provide variations of a given functional with respect to change of certain material properties, and not a domain topology. Recently, topological asymptotic analysis has been also applied for image processing purposes by Auroux et al. [8, 9] and by Muszkieta [37].

In order to apply the theory of topological asymptotic expansions, it is necessary that the functional to be minimized depends solely on the set K. This can be achieved in the case of the Mumford–Shah functional by noting that the minimizing pair (u_0, K_0) is uniquely determined by either of the components u_0 and K_0: The set K_0 coincides with the jump set S_{u_0} of the function u_0. Conversely, u_0 can be computed from K_0 by solving the partial differential equation

$$\begin{aligned} u - \alpha \Delta u &= f \quad \text{in } \Omega\,, \\ \frac{\partial u}{\partial n} &= 0 \quad \text{in } \partial(\Omega \setminus K_0)\,. \end{aligned} \tag{1.2}$$

Now consider the functional

$$J(K) = F\big(u(K), K\big)\,,$$

where $u(K)$ denotes the solution of (1.2) with K_0 replaced by K. Then the pair $(u(K_0), K_0)$ minimizes the functional F, if and only if $J(K_0) \le J(K)$ for all $K \subset \Omega$.

The idea is now to use a gradient descent like approach to the minimization of the functional J. Starting from an initial guess K of the edge set (for instance $K = \emptyset$), one adds to K those points, whose inclusion would lead to a near to maximal decrease of the cost functional J. More precisely, one adjoins to the set K small balls of radius $\varepsilon > 0$ centered at the points $x \in \Omega \setminus K$ and tries to compute an asymptotic expansion of the form

$$J\big(K \cup B_\varepsilon(x)\big) - J(K) = c(\varepsilon) G_K(x) + o\big(c(\varepsilon)\big)$$

for some functions $c\colon \mathbb{R}_{>0} \to \mathbb{R}_{>0}$ and $G_K\colon \Omega \setminus K \to \mathbb{R}$. Those $x \in \Omega \setminus K$ for which G_K attains the largest negative values are then added to the set K. This process is iterated, until the function G_K becomes non-negative everywhere.

In the case of the Mumford–Shah functional, this procedure cannot applied directly, because the functional is infinite whenever K contains a set of positive Lebesgue measure. We therefore propose to use a different, though related, functional for the computation of the asymptotic expansion, which is based on an approximation of the one-dimensional Hausdorff measure: The number of balls of radius $\varepsilon > 0$ that are required to cover a set K, multiplied with 2ε, provides a good estimate of $\mathcal{H}^1(K)$ for ε small enough and K sufficiently regular. In the following, we introduce this approximating functional $J_{\varepsilon,\kappa}$. However, because we later prove the Γ-convergence of this functional to F, it is necessary to let $J_{\varepsilon,\kappa}$ depend on two functions, the function u and a piecewise constant edge indicator function v:

For each finite set $Y \subset \mathbb{R}^2$ and each $0 < \kappa < 1$ we define the function $v_{Y,\kappa}\colon \Omega \to \mathbb{R}$ by

$$v_{Y,\kappa}(x) := \begin{cases} \kappa & \text{if } x \in \bigcup_{y \in Y} B_\varepsilon(y)\,, \\ 1 & \text{else.} \end{cases}$$

For every $v \in L^2(\Omega)$ we define

$$m_{\varepsilon,\kappa}(v) := \inf\big\{\mathcal{H}^0(Y) : Y \subset \mathbb{R}^2,\ v = v_{Y,\kappa}\big\}\ .$$

Here we set $m_{\varepsilon,\kappa}(v) := +\infty$, if $v \ne v_{Y,\kappa}$ for any $Y \subset \mathbb{R}^2$. Note that, for a given function v, there might exist different sets $Y \subset \mathbb{R}^2$ such that $v = v_{Y,\kappa}$.

Finally, we introduce the family of functionals $J_{\varepsilon,\kappa}\colon L^2(\Omega) \times L^2(\Omega) \to \mathbb{R}$, defined by

$$J_{\varepsilon,\kappa}(u, v) := \frac{1}{2}\int_\Omega (u - f)^2\,dx + \frac{\alpha}{2}\int_\Omega v|\nabla u|^2\,dx + 2\beta\varepsilon\, m_{\varepsilon,\kappa}(v) \tag{1.3}$$

if $u \in H^1(\Omega)$, and $J_\varepsilon(u, v) := +\infty$ otherwise.

In Section 2, we derive an approximation of the functional $J_{\varepsilon,\kappa}$ that allows us to compute an approximation of the minimizer using the idea of topological asymptotic expansions. We show that

$$J_{\varepsilon,\kappa}(u, v_{Y\cup\{y\},\kappa}) - J_{\varepsilon,\kappa}(u, v_{Y,\kappa}) \approx -\varepsilon^2 \pi\alpha \frac{1-\kappa}{1+\kappa} |\nabla u(y)|^2 + 2\varepsilon\beta \; .$$

(cf. Theorem 2.1).

In Section 3, we show that the functional $J_{\varepsilon,\kappa}$ is indeed an approximation of the Mumford–Shah functional in the sense of Γ-convergence. More precisely, if we choose $\kappa = \kappa(\varepsilon)$ in such a way that $\kappa(\varepsilon) = o(\varepsilon)$ as $\varepsilon \to 0$, then

$$F = \Gamma\text{-}\lim_{\varepsilon\to 0} J_{\varepsilon,\kappa(\varepsilon)} \; .$$

In particular, this implies that the minimizers of $J_{\varepsilon,\kappa(\varepsilon)}$ converge to a minimizer of F as ε tends to zero. The adopted proof is based on the methods used for proving the Γ-convergence of the Ambrosio–Tortorelli approximation of F (see [3, 11]). Finally, in the Section 4, we propose an algorithm to minimize the functional $J_{\varepsilon,\kappa}$. We compare numerical results obtained with this algorithm with results obtained by minimization of the Ambrosio and Tortorelli model [3].

2 Topological Asymptotic Analysis

In this section, we derive the topological asymptotic expansion of the functional $J_{\varepsilon,\kappa}$ defined in (1.3). This expansion will allow us to find its approximate minimizers. We assume that $\Omega \subset \mathbb{R}^2$ is a bounded and open Lipschitz domain and that $f \in H^1(\Omega)$ is given. We define the functional $G\colon L^2(\Omega) \times L^2(\Omega) \to [0, +\infty]$,

$$G(\tilde u, \tilde v) := \frac{1}{2}\int_\Omega (\tilde u - f)^2 \, dx + \frac{\alpha}{2}\int_\Omega \tilde v |\nabla \tilde u|^2 \, dx \; .$$

Now assume that K is an open subset of Ω and $0 < \kappa < 1$ satisfying $\alpha\kappa < 1$. We define the function $v\colon \Omega \to \mathbb{R}$ by

$$v(x) = \begin{cases} \kappa & \text{if } x \in K, \\ 1 & \text{else.} \end{cases}$$

Using standard methods of variational calculus one can show that the mapping $\tilde u \mapsto G(\tilde u, v)$ attains a unique minimizer in $H^1(\Omega)$, which we denote by u.

The main result of this section is stated in the following theorem:

Theorem 2.1 *Let $K \subset \Omega$. For $\hat y \in \Omega \setminus \bar K$ and $\varepsilon > 0$ define the functions*

$$v^{(\hat y,\varepsilon)}(x) := \begin{cases} \kappa & \textit{if } x \in K \cup B_\varepsilon(\hat y), \\ 1 & \textit{else,} \end{cases}$$

and

$$u^{(\hat y,\varepsilon)} := \operatorname*{arg\,min}_{\tilde u \in H^1(\Omega)} G(\tilde u, v^{(\hat y,\varepsilon)}) \, .$$

Then for all compact subsets $L \subset \Omega \setminus \bar K$ we have

$$\lim_{\varepsilon\to 0} \sup_{\hat y\in L} \frac{1}{\varepsilon^2} \left| G(u^{(\hat y,\varepsilon)}, v^{(\hat y,\varepsilon)}) - G(u,v) + \alpha\varepsilon^2\pi \frac{1-\kappa}{1+\kappa} |\nabla u(\hat y)|^2 \right| = 0 \; .$$

For the remaining part of this section we assume that the compact set $L \subset \Omega \setminus \bar{K}$ is fixed. Moreover we define

$$\delta := \frac{1}{3}\operatorname{dist}(L, \partial\Omega \cup \bar{K}) = \frac{1}{3}\min\bigl\{|x-y| : x \in L,\ y \in \partial\Omega \cup \bar{K}\bigr\}\ .$$

In addition, we consider the set

$$\hat{L} := L + \bar{B}_\delta(0) := \bigl\{x + y : x \in L,\ |y| \le \delta\bigr\}\ .$$

Before we give the proof of the Theorem 2.1, we need to introduce some auxiliary results. First, we recall that the assumptions that u and $u^{(\hat{y},\varepsilon)}$ are minimizers of $G(\cdot, v)$ and $G(\cdot, v^{(\hat{y},\varepsilon)})$, respectively, imply that

$$\begin{aligned} \int_\Omega (u^{(\hat{y},\varepsilon)} - f)\varphi + \alpha v^{(\hat{y},\varepsilon)} \langle \nabla u^{(\hat{y},\varepsilon)}, \nabla\varphi\rangle\, dx &= 0\,, \\ \int_\Omega (u - f)\varphi + \alpha v \langle \nabla u, \nabla\varphi\rangle\, dx &= 0\,, \end{aligned} \tag{2.1}$$

for all $\varphi \in H^1(\Omega)$.

We first we need a regularity result for the function u.

Lemma 2.1 *The function u satisfies*

$$u \in C^1_{loc}(\Omega \setminus \bar{K})\ .$$

Moreover, there exists a constant C_1 only depending on L and K such that

$$\|\nabla u\|_{L^\infty(\hat{L})} \le C_1 \|f\|_{H^1(\Omega)}\ .$$

Proof. Because by assumption $f \in H^1(\Omega)$ and the function v is constant on $\Omega \setminus K$, it follows from standard theorems on the regularity of solutions of elliptic equations that $u \in H^3_{\mathrm{loc}}(\Omega \setminus \bar{K})$ (see [31, Thm. 8.10]). Moreover, there exists a constant $\tilde{C}$ only depending on $\hat{L}$ (and therefore on L and K) such that $\|u\|_{H^3(\hat{L})} \le \tilde{C}\|f\|_{H^1(\Omega)}$. Thus the Sobolev embedding theorem [1, Thm. 5.4] implies the assertion. $\square$

As a second step, we need H^1-norm and L^2-norm estimates of the difference $u^{(\hat{y},\varepsilon)} - u$. First we show that the H^1-norm of the difference $u^{(\hat{y},\varepsilon)} - u$ on the whole domain Ω is of order ε.

Lemma 2.2 *There exists a constant $C_2 > 0$ only depending on L, κ, Ω, and K such that for all $\varepsilon > 0$ with $\varepsilon < \delta$ and $\hat{y} \in L$ the estimate*

$$\|u^{(\hat{y},\varepsilon)} - u\|_{H^1(\Omega)} \le C_2 \|f\|_{H^1(\Omega)}\varepsilon$$

holds.

Proof. Computing the difference between the two equations in (2.1) and using the definition of $v^{(\hat{y},\varepsilon)}$, we obtain that

$$\begin{aligned} \int_\Omega (u^{(\hat{y},\varepsilon)} - u)\varphi\, dx + \alpha \int_\Omega v^{(\hat{y},\varepsilon)} \langle \nabla(u^{(\hat{y},\varepsilon)} - u), \nabla\varphi\rangle\, dx& \\ = \alpha(1-\kappa)\int_{B_\varepsilon(\hat{y})} \langle \nabla u, \nabla\varphi\rangle\, dx& \end{aligned}$$

for all $\varphi \in H^1(\Omega)$. In particular, it follows with $\varphi = u^{(\hat{y},\varepsilon)} - u$ that

$$\begin{aligned}\alpha\kappa \|u^{(\hat{y},\varepsilon)} - u\|^2_{H^1(\Omega)} &\le \alpha(1-\kappa) \int_{B_\varepsilon(\hat{y})} \langle \nabla u, \nabla(u^{(\hat{y},\varepsilon)} - u)\rangle \, dx \\ &\le \alpha(1-\kappa) \|\nabla u\|_{L^2(B_\varepsilon(\hat{y}))} \|u^{(\hat{y},\varepsilon)} - u\|_{H^1(\Omega)} \, .\end{aligned}$$

Moreover Lemma 2.1 implies that

$$\|\nabla u\|_{L^2(B_\varepsilon(\hat{y}))} \le \|\nabla u\|_{L^\infty(B_\varepsilon(\hat{y}))} \sqrt{\pi}\varepsilon \le \|\nabla u\|_{L^\infty(\hat{L})} \sqrt{\pi}\varepsilon \le C_1 \sqrt{\pi} \|f\|_{H^1(\Omega)} \varepsilon \, . \tag{2.2}$$

Setting $C_2 := (1-\kappa)\sqrt{\pi} C_1/\kappa$, the assertion follows. □

Lemma 2.3 *There exists a constant C_3 only depending on L, κ, Ω, K, and $\|f\|_{H^1(\Omega)}$ such that*

$$\|u^{(\hat{y},\varepsilon)} - u\|_{L^2(\Omega)} \le C_3 \varepsilon^{3/2}$$

for every $\hat{y} \in L$ and $0 < \varepsilon < \delta$.

Proof. Let $g \in H^1(\Omega)$ satisfy $\|g\|_{H^1(\Omega)} \le 1$ and assume that $w^{(\hat{y},\varepsilon)}$ and w are the unique solutions to the equations (2.1) but with given g instead of f, that is

$$\begin{aligned}\int_\Omega (w^{(\hat{y},\varepsilon)} - g)\varphi + \alpha v^{(\hat{y},\varepsilon)} \langle \nabla w^{(\hat{y},\varepsilon)}, \nabla\varphi\rangle \, dx &= 0 \, , \\ \int_\Omega (w - g)\varphi + \alpha v \langle \nabla w, \nabla\varphi\rangle \, dx &= 0 \, ,\end{aligned} \tag{2.3}$$

for all $\varphi \in H^1(\Omega)$.

Taking $\varphi = w^{(\hat{y},\varepsilon)}$ and $\varphi = w$ in the first and the second equation in (2.1), respectively, and next subtracting these equations from the corresponding equations in (2.3) with $\varphi = u^{(\hat{y},\varepsilon)}$ and $\varphi = u$, we obtain

$$\begin{aligned}\int_\Omega w^{(\hat{y},\varepsilon)} \, f \, dx - \int_\Omega u^{(\hat{y},\varepsilon)} \, g \, dx &= 0 \, , \\ \int_\Omega w \, f \, dx - \int_\Omega u \, g \, dx &= 0 \, .\end{aligned}$$

In particular,

$$\int_\Omega (u^{(\hat{y},\varepsilon)} - u) \, g \, dx = \int_\Omega (w^{(\hat{y},\varepsilon)} - w) \, f \, dx \, .$$

Next, we note that

$$\begin{aligned}\int_\Omega (u^{(\hat{y},\varepsilon)} - u) \, g \, dx &= \frac{1}{2} \left(\int_\Omega (u^{(\hat{y},\varepsilon)} - u) \, g \, dx + \int_\Omega (w^{(\hat{y},\varepsilon)} - w) \, f \, dx \right) \\ &= \frac{1}{2} \int_\Omega \Big((u^{(\hat{y},\varepsilon)} + w^{(\hat{y},\varepsilon)}) - (u + w) \Big) \, (f + g) \, dx \\ &\qquad - \frac{1}{2} \int_\Omega (u^{(\hat{y},\varepsilon)} - u) \, f \, dx - \frac{1}{2} \int_\Omega (w^{(\hat{y},\varepsilon)} - w) \, g \, dx \, .\end{aligned} \tag{2.4}$$

Computing the difference of the two equations in (2.1) with $\varphi = u$ and $\varphi = u^{(\hat{y},\varepsilon)}$, respectively, we obtain

$$\begin{aligned}\int_\Omega (u^{(\hat{y},\varepsilon)} - u)\, f\, dx &= \alpha \left(\int_\Omega v \langle \nabla u, \nabla u^{(\hat{y},\varepsilon)} \rangle \, dx - \int_\Omega v^{(\hat{y},\varepsilon)} \langle \nabla u, \nabla u^{(\hat{y},\varepsilon)} \rangle \, dx \right) \\ &= \alpha(1-\kappa) \int_{B_\varepsilon(\hat{y})} \langle \nabla u, \nabla u^{(\hat{y},\varepsilon)} \rangle \, dx \\ &= \alpha(1-\kappa) \left(\int_{B_\varepsilon(\hat{y})} \langle \nabla u, \nabla (u^{(\hat{y},\varepsilon)} - u) \rangle \, dx + \int_{B_\varepsilon(\hat{y})} |\nabla u|^2 \, dx \right)\end{aligned}$$

Application of the Cauchy-Schwarz inequality to the above formula, and next, the estimate (2.2) and Lemma 2.2 yields the inequality

$$\begin{aligned}&\Big| \int_\Omega (u^{(\hat{y},\varepsilon)} - u)\, f\, dx \Big| \\ &\quad \le \alpha(1-\kappa) \left(\|\nabla u\|_{L^2(B_\varepsilon(\hat{y}))} \|u^{(\hat{y},\varepsilon)} - u\|_{H^1(\Omega)} + \|\nabla u\|^2_{L^2(B_\varepsilon(\hat{y}))} \right) \\ &\quad \le \alpha(1-\kappa) \left(\sqrt{\pi} C_1 C_2 + \pi C_1^2 \right) \|f\|^2_{H^1(\Omega)} \varepsilon^2\end{aligned} \tag{2.5}$$

In a similar manner, using the assumption that $\|g\|_{H^1(\Omega)} = 1$, we can show that

$$\Big| \int_\Omega (w^{(\hat{y},\varepsilon)} - w) g\, dx \Big| \le \alpha(1-\kappa) \left(\sqrt{\pi} C_1 C_2 + \pi C_1^2 \right) \varepsilon^2 \,, \tag{2.6}$$

and

$$\begin{aligned}&\Big| \int_\Omega \big((u^{(\hat{y},\varepsilon)} + w^{(\hat{y},\varepsilon)}) - (u + w) \big)(f + g)\, dx \Big| \\ &\qquad\qquad \le \alpha(1-\kappa) \left(\sqrt{\pi} C_1 C_2 + \pi C_1^2 \right) 2(\|f\|^2_{H^1(\Omega)} + 1) \varepsilon^2 \,.\end{aligned} \tag{2.7}$$

Finally, combining (2.5), (2.6) and (2.7) with (2.4), we obtain

$$\begin{aligned}&\Big| \int_\Omega (u^{(\hat{y},\varepsilon)} - u) g\, dx \Big| \\ &\qquad \le \frac{3\alpha(1-\kappa)}{2} \left(\sqrt{\pi} C_1 C_2 + \pi C_1^2 \right) (\|f\|^2_{H^1(\Omega)} + 1) \varepsilon^2 \,.\end{aligned}$$

Therefore, we have

$$\begin{aligned}\|u^{(\hat{y},\varepsilon)} - u\|_{H^{-1}(\Omega)} &:= \sup \Big\{ \Big| \int_\Omega (u^{(\hat{y},\varepsilon)} - u) g\, dx \Big| : g \in H^1(\Omega),\ \|g\|_{H^1(\Omega)} \le 1 \Big\} \\ &\le c(\|f\|^2_{H^1(\Omega)} + 1) \varepsilon^2\end{aligned} \tag{2.8}$$

with $c = 3\alpha(1-\kappa) \left(\sqrt{\pi} C_1 C_2 + \pi C_1^2 \right) /2$. Now, estimates from the theory of Hilbert scales (see [33, Thm. 9.4]), Lemma 2.2, and (2.8) imply that

$$\begin{aligned}\|u^{(\hat{y},\varepsilon)} - u\|^2_{L^2(\Omega)} &\le \|u^{(\hat{y},\varepsilon)} - u\|_{H^{-1}(\Omega)} \|u^{(\hat{y},\varepsilon)} - u\|_{H^1(\Omega)} \\ &\le c(\|f\|^2_{H^1(\Omega)} + 1) \varepsilon^2 \, C_2 \|f\|_{H^1(\Omega)} \varepsilon \,.\end{aligned}$$

Therefore, the desired estimate holds with $C_3 = \sqrt{c(\|f\|^2_{H^1(\Omega)} + 1) C_2 \|f\|_{H^1(\Omega)}}$. □

In the next step, we need to derive an estimate for the function $u^{(\hat{y},\varepsilon)}$ on the boundary $\partial B_\varepsilon(\hat{y})$. To do this, we follow Vogelius et al. [43], where such an estimate has been derived for the solution of the homogeneous Helmholtz equation with Dirichlet boundary conditions.

We first introduce the Green function corresponding to the equation $u - \alpha\Delta u = f$ on $\Omega \setminus \bar{K}$ with Neumann boundary conditions. That is, the function $N(\cdot, y)$ solves the problem

$$\begin{cases} N(x,y) - \alpha\Delta_x N(x,y) = \delta_y(x) & x \in \Omega \setminus \bar{K}\,, \\ \dfrac{\partial N}{\partial n}(x,y) = 0 & x \in \partial(\Omega \setminus \bar{K})\,, \end{cases}$$

for $y \in \Omega$. We note that $N(\cdot, y)$ can be written as the sum of the fundamental solution corresponding to the equation $u - \alpha\Delta u = \delta_y$, denoted by $\Gamma(\cdot, y)$, and a corrector function $h(\cdot, y)$, which is chosen in such a way that the normal derivative of $N(\cdot, y)$ vanishes on the boundary $\partial(\Omega \setminus \bar{K})$. The function $\Gamma(\cdot, y)$ is given by

$$\Gamma(\cdot, y) = \frac{1}{2\pi} K_0\left(\frac{1}{\sqrt{\alpha}}|x-y|\right)$$

for all x, $y \in \mathbb{R}^2$, such that $x \neq y$. Here K_0 denotes the modified Bessel function of the second kind (see, e.g., [38, p. 490]). Furthermore, the function K_0 has an asymptotic expansion of the form

$$K_0(z) = -\ln z + \ln 2 - \gamma + O(z^2|\ln z|)$$

for $z \to 0$, where γ denotes the Euler-Mascheroni constant (see, e.g., [42, Ch. 51]). Therefore, we conclude that $\Gamma(\cdot, y)$ can be approximated as

$$\Gamma(x,y) = \frac{1}{2\pi}\Big(-\ln(|x-y|) + \frac{1}{2}\ln\alpha + \ln 2 - \gamma\Big) + O(|x-y|^2|\ln|x-y||)\,, \quad (2.9)$$

when $|x - y| \to 0$. Moreover, we observe that $\Gamma(\cdot, y)$ has the same singular behavior as the fundamental solution of the Laplace equation

$$\Phi(x,y) = -\frac{1}{2\pi}\ln(|x-y|)$$

defined for all x, $y \in \mathbb{R}^2$, such that $x \neq y$. We need the approximation (2.9) in order to be able to apply standard methods of potential theory to derive an estimation for the function $u^{(\hat{y},\varepsilon)}$ on $\partial B_\varepsilon(\hat{y})$. Such way of proceeding is common when dealing with problems of this kind (see, e.g., Colton and Kress [20, 21]).

Lemma 2.4 *There exists a constant C_4 only depending on L, K, and Ω, such that for every point $\hat{y} \in L$, $0 < \varepsilon < \delta$, and $y \in \Omega$ satisfying $\varepsilon < |y - \hat{y}| < 2\varepsilon$ the estimate*

$$\|N(\cdot, y)\|_{L^2(B_\varepsilon)} \leq C_4 \varepsilon |\ln \varepsilon|$$

holds.

Proof. Since by elliptic regularity $h \in C^\infty(\Omega \setminus K, \Omega \setminus \bar{K})$ and

$$\operatorname{dist}(\hat{y}, \Omega \setminus K) \geq 3\delta > 3\varepsilon\,,$$

we have

$$\|h(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))} \le \sqrt{\pi}\|h(\cdot,y)\|_{L^\infty(\Omega\setminus\bar K)}\varepsilon \le \sqrt{\pi}c_1\varepsilon$$

with $c_1 > 0$ only depending on L, K, and Ω. The estimate $\|\Phi(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))} \le c_2\varepsilon|\ln\varepsilon|$ for some c_2 only depending on ε and for all y satisfying $\varepsilon < |y-\hat y| < 2\varepsilon$ can be easily derived in the polar coordinate system. Therefore, from the Minkowski inequality and (2.9) we get that

$$\begin{aligned}\|N(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))} &\le \|h(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))} + \|\Gamma(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))}\\ &\le \sqrt{\pi}c_1\varepsilon + c_2\varepsilon|\ln\varepsilon| \le C_4\varepsilon|\ln\varepsilon|\end{aligned}$$

with C_4 chosen slightly larger than c_2. □

Lemma 2.5 *There exists a constant C_5 only depending on L, K, Ω κ, α, and f, such that for every point $\hat y \in L$, $0 < \varepsilon < \delta$, and $y \in \Omega$ satisfying $\varepsilon < |y-\hat y| < 2\varepsilon$ the estimate*

$$\left|u^{(\hat y,\varepsilon)}(y) - u(y) - \alpha(1-\kappa)\int_{\partial B_\varepsilon(\hat y)} u^{(\hat y,\varepsilon)}(x)\frac{\partial N}{\partial n}(x,y)\,ds(x)\right| \le C_5\varepsilon^{3/2}$$

holds.

Proof. Computing the difference of the integral representation formulas of the functions u and $u^{(\hat y,\varepsilon)}$, we obtain

$$\begin{aligned}&u^{(\hat y,\varepsilon)}(y) - u(y) =\\ &\alpha\int_{\partial B_\varepsilon(\hat y)}\left(u^{(\hat y,\varepsilon)}(x)\frac{\partial N}{\partial n}(x,y) - N(x,y)\frac{\partial u^{(\hat y,\varepsilon)^+}}{\partial n}(x)\right)ds(x)\\ &-\alpha\int_{\partial K} N(x,y)\frac{\partial}{\partial n}\left(u^{(\hat y,\varepsilon)^+}(x) - u^+(x)\right)ds(x)\\ &-\alpha\int_{B_\varepsilon(\hat y)} N(x,y)\,f(x)\,dx\,.\end{aligned}\tag{2.10}$$

for all $y \in \Omega\setminus(\bar K\cup\bar B_\varepsilon(\hat y))$.

Using the transmission condition that $u^{(\hat y,\varepsilon)}$ satisfies on $\partial B_\varepsilon(\hat y)$, the Green formula, and that

$$N(x,y) - \alpha\Delta N(x,y) = 0 \tag{2.11}$$

for $x \in B_\varepsilon(\hat y)$ and $y \in \Omega\setminus(\bar K\cup\bar B_\varepsilon(\hat y))$, we have

$$\begin{aligned}&\alpha\int_{\partial B_\varepsilon(\hat y)} N(x,y)\frac{\partial u^{(\hat y,\varepsilon)^+}}{\partial n}(x)\,ds(x) = \alpha\kappa\int_{\partial B_\varepsilon(\hat y)} N(x,y)\frac{\partial u^{(\hat y,\varepsilon)^-}}{\partial n}(x)\,ds(x)\\ &= \alpha\kappa\int_{B_\varepsilon(\hat y)} N(x,y)\,\Delta u^{(\hat y,\varepsilon)}(x)\,dx + \alpha\kappa\int_{B_\varepsilon(\hat y)}\left\langle\nabla N(x,y),\nabla u^{(\hat y,\varepsilon)}(x)\right\rangle dx\\ &= (1-\kappa)\int_{B_\varepsilon(\hat y)} N(x,y)u^{(\hat y,\varepsilon)}(x)\,dx - \int_{B_\varepsilon(\hat y)} N(x,y)f(x)\,dx\\ &\quad + \alpha\kappa\int_{\partial B_\varepsilon(\hat y)} u^{(\hat y,\varepsilon)}(x)\frac{\partial N}{\partial n}(x,y)\,ds(x)\,.\end{aligned}\tag{2.12}$$

Using similar arguments as above and applying the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned}
&-\alpha \int_{\partial K} N(x,y)\frac{\partial}{\partial n}\big({u^{(\hat{y},\varepsilon)}}^{+}(x) - u^{+}(x)\big)\,ds(x)\\
&\qquad\qquad = -\alpha\kappa \int_{\partial K} N(x,y)\frac{\partial}{\partial n}\big({u^{(\hat{y},\varepsilon)}}^{-}(x) - u^{-}(x)\big)\,ds(x)\\
&\qquad\qquad = \alpha(1-\kappa)\int_K \big(u^{(\hat{y},\varepsilon)}(x) - u(x)\big)N(x,y)\,dx\\
&\qquad\qquad \le \alpha(1-\kappa)\|u^{(\hat{y},\varepsilon)} - u\|_{L^2(K)}\|N(\cdot,y)\|_{L^2(K)}\,.
\end{aligned} \tag{2.13}$$

Because $\operatorname{dist}(y,K) \ge \operatorname{dist}(\hat{y},K) - 2\varepsilon > \delta$, there exists some $c > 0$ such that $\|N(\cdot,y)\|_{L^2(K)} < c$. Application of Lemma 2.3 therefore yields

$$\|u^{(\hat{y},\varepsilon)} - u\|_{L^2(K)}\|N(\cdot,y)\|_{L^2(K)} \le c\|u^{(\hat{y},\varepsilon)} - u\|_{L^2(\Omega)} \le C_3 c\,\varepsilon^{3/2}\,. \tag{2.14}$$

Taking into account (2.12), (2.13), and (2.14) in (2.10), we obtain

$$\begin{aligned}
u^{(\hat{y},\varepsilon)}(y) - u(y) =& \alpha(1-\kappa)\int_{\partial B_\varepsilon(\hat{y})} u^{(\hat{y},\varepsilon)}(x)\frac{\partial N}{\partial n}(x,y)\,ds(x)\\
&- (1-\kappa)\int_{B_\varepsilon(\hat{y})} u^{(\hat{y},\varepsilon)}(x)N(x,y)\,dx\\
&+ (1-\alpha)\int_{B_\varepsilon(\hat{y})} f(x)N(x,y)\,dx + O(\varepsilon^{3/2})\,,
\end{aligned}$$

the last term being bounded by $\alpha(1-\kappa)C_3 c\,\varepsilon^{3/2}$. To estimate the remaining integrals, we note that

$$\|u^{(\hat{y},\varepsilon)}\|_{L^2(B_\varepsilon(\hat{y}))} \le \sqrt{\pi}\|f\|_{L^\infty(\Omega)}\,\varepsilon\,. \tag{2.15}$$

Next, using the Cauchy–Schwarz inequality and Lemma 2.4 we get

$$\begin{aligned}
\int_{B_\varepsilon(\hat{y})} u^{(\hat{y},\varepsilon)}(x)N(x,y)\,dx &\le \|u^{(\hat{y},\varepsilon)}\|_{L^2(B_\varepsilon(\hat{y}))}\|N(\cdot,y)\|_{L^2(B_\varepsilon(\hat{y}))}\\
&\le \sqrt{\pi}\|f\|_{L^\infty(\Omega)}C_4\,\varepsilon^2|\ln\varepsilon|\,.
\end{aligned}$$

In a similar way, we show that

$$\begin{aligned}
\int_{B_\varepsilon(\hat{y})} f(x)N(x,y)\,dx &\le \|f\|_{L^2(B_\varepsilon(\hat{y}))}\|N(\cdot,y)\|_{L^2(B_\varepsilon(\hat{y}))}\\
&\le \sqrt{\pi}\|f\|_{L^\infty(\Omega)}C_4\,\varepsilon^2|\ln\varepsilon|\,.
\end{aligned}$$

Therefore, we obtain the desired estimate with C_5 slightly larger than $\alpha(1-\kappa)C_3c$. $\square$

Lemma 2.6 *There exists a constant C_6 only depending on L, K, κ, α, and f, such that for every point $\hat{y} \in L$, $0 < \varepsilon < \delta$, and $y \in \Omega$ satisfying $\varepsilon < |y - \hat{y}| < 2\varepsilon$ the estimate*

$$\Big|\int_{\partial B_\varepsilon(\hat{y})} u^{(\hat{y},\varepsilon)}(x)\frac{\partial N}{\partial n}(x,y)\,ds(x) - \int_{\partial B_\varepsilon(\hat{y})} u^{(\hat{y},\varepsilon)}(x)\frac{\partial \Phi}{\partial n}(x,y)\,ds(x)\Big| \le C_6\varepsilon^2|\ln\varepsilon|$$

holds.

Proof. Denote $H(x,y) := h(x,y) + \Gamma(x,y) - \Phi(x,y)$. Then $N(x,y) = \Phi(x,y) + H(x,y)$ and, using (2.11), we obtain

$$N(x,y) = \Delta H(x,y)\,. \tag{2.16}$$

Application of the Green formula, (2.16), and the Cauchy-Schwarz inequality yields

$$\begin{aligned}
&\Big|\int_{\partial B_\varepsilon(\hat y)} u^{(\hat y,\varepsilon)}(x)\frac{\partial H}{\partial n}(x,y)\,ds(x)\Big| \\
&\qquad = \Big|\int_{B_\varepsilon(\hat y)} u^{(\hat y,\varepsilon)}(x)\Delta H(x,y)\,dx + \int_{B_\varepsilon(\hat y)} \big\langle \nabla u^{(\hat y,\varepsilon)}(x), \nabla H(x,y)\big\rangle\,dx\Big| \\
&\qquad = \Big|\int_{B_\varepsilon(\hat y)} u^{(\hat y,\varepsilon)}(x)N(x,y)\,dx + \int_{B_\varepsilon(\hat y)} \big\langle \nabla u^{(\hat y,\varepsilon)}(x), \nabla H(x,y)\big\rangle\,dx\Big| \\
&\qquad \le \|u^{(\hat y,\varepsilon)}\|_{L^2(B_\varepsilon(\hat y))}\|N(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))} \\
&\qquad\quad + \|\nabla(u^{(\hat y,\varepsilon)} - u)\|_{L^2(B_\varepsilon(\hat y))}\,\|\nabla H(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))} \\
&\qquad\quad + \|\nabla u\|_{L^2(B_\varepsilon(\hat y))}\,\|\nabla H(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))}
\end{aligned}$$

Using (2.15) and Lemmas 2.2 and 2.4, it follows that

$$\begin{aligned}
&\Big|\int_{\partial B_\varepsilon(\hat y)} u^{(\hat y,\varepsilon)}(x)\frac{\partial H}{\partial n}(x,y)\,ds(x)\Big| \\
&\le C_4\sqrt{\pi}\|f\|_{L^\infty(\Omega)}\varepsilon^2|\ln\varepsilon| + (C_2 + C_1\sqrt{\pi})\|f\|_{H^1(\Omega)}\|\nabla H(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))}\varepsilon
\end{aligned} \tag{2.17}$$

Using the explicit form of $\Gamma(\cdot,y)$ and $\Phi(x,y)$, and next the Taylor expansion of the function K_0 (see [42, Ch. 51]), we can show that

$$|\nabla\Gamma(x,y) - \nabla\Phi(x,y)| \le c\,\varepsilon|\ln\varepsilon|$$

for some $c > 0$ only depending on δ and all $x \in B_\varepsilon(\hat y)$ and y satisfying $\varepsilon < |y - \hat y| < 2\varepsilon$. Thus

$$\|\nabla H(\cdot,y)\|_{L^2(B_\varepsilon(\hat y))} \le (c\varepsilon|\ln\varepsilon| + \|h\|_{L^\infty(B_\varepsilon(\hat y))})\sqrt{\pi}\varepsilon$$

Thus we obtain the required estimate from (2.17) by choosing C_6 slightly larger than $C_4\sqrt{\pi}\|f\|_{L^\infty(\Omega)}$. □

From Lemma 2.5 and Lemma 2.6 and using the jump formula for the double layer potential (see e.g. Kress [34, p. 67]), we have

$$\begin{aligned}
&\Big|\frac{1}{2}\alpha(1+\kappa)u^{(\hat y,\varepsilon)}(y) - u(y) \\
&\qquad -\alpha(1-\kappa)\int_{\partial B_\varepsilon(\hat y)} u^{(\hat y,\varepsilon)}(x)\frac{\partial\Phi}{\partial n}(x,y)\,ds(x)\Big| \le C_7\,\varepsilon^{3/2}
\end{aligned} \tag{2.18}$$

for $y \in \partial B_\varepsilon(\hat y)$, with $C_7 > C_5$.

Now we introduce the auxiliary vector valued function $\phi\colon \mathbb{R}^2 \to \mathbb{R}\times\mathbb{R}$ that solves the problem

$$\begin{cases} \Delta\phi(x) = 0 & x \in \mathbb{R}^2 \setminus \bar{B}(0) \text{ or } x \in B(0)\,, \\ \phi^+(y) = \phi^-(x) & x \in \partial B(0)\,, \\ \dfrac{\partial \phi^+}{\partial n}(x) - \kappa \dfrac{\partial \phi^-}{\partial n}(x) = -\kappa n(x) & x \in \partial B(0)\,, \\ \lim\limits_{|x|\to\infty} \phi(x) = 0\,. \end{cases} \tag{2.19}$$

Here $B(0)$ denotes the unit ball in $\mathbb{R}^2$ of a center in 0 and n is the unit normal vector exterior to the boundary $\partial B(0)$. The existence and uniqueness of a solution to the problem (2.19) can be proved using single layer potentials with suitably chosen densities. For details, see Ammari and Kang [4] or Cedio-Fengya et al. [14].

Solving the problem (2.19) using standard methods of potential theory, we obtain the explicit form of ϕ, which reads

$$\phi(x) = \frac{\kappa}{\kappa+1} x \quad \text{and} \quad \phi(x) = \frac{\kappa}{\kappa+1}\frac{x}{|x|^2} \tag{2.20}$$

for all $x \in B(0)$ and $x \in \mathbb{R}^2 \setminus \bar{B}(0)$, respectively.

The result on asymptotic expansion of the function $u^{(\hat{y},\varepsilon)}$ on the boundary $\partial B_\varepsilon(\hat{y})$ is stated in the following Lemma:

Lemma 2.7 *For every point $\hat{y} \in L$, $0 < \varepsilon < \delta$, and $y \in \partial B_\varepsilon(\hat{y})$ the estimate*

$$\left| u^{(\hat{y},\varepsilon)}(y) - u(y) - \varepsilon \left(\frac{1}{\kappa} - 1\right) \langle \phi(y/\varepsilon), \nabla u(\hat{y})\rangle \right| \le C_7 \varepsilon^{3/2}$$

holds, where the constant C_7 is as in (2.18).

Proof. The proof of this lemma can be proved starting from the formula (2.18) in the same way as in Vogelius and Volkov [43, Prop. 3]. □

Using all the above results, we can now prove Theorem 2.1.

2.1 Proof of Theorem 2.1

Using (2.1) with $\varphi = u^{(\hat{y},\varepsilon)}$ and $\varphi = u$, we obtain that

$$G(u^{(\hat{y},\varepsilon)}, v^{(\hat{y},\varepsilon)}) - G(u,v) = -\frac{1}{2}\int_\Omega f(u^{(\hat{y},\varepsilon)} - u)\,dx\,. \tag{2.21}$$

Again using (2.1), it follows that

$$\begin{aligned} -\frac{1}{2}\int_\Omega & f(u^{(\hat{y},\varepsilon)} - u)\,dx \\ &= \frac{1}{2}\int_\Omega v^{(\hat{y},\varepsilon)} \langle \nabla u, \nabla u^{(\hat{y},\varepsilon)}\rangle\,dx - \frac{1}{2}\int_\Omega v \langle \nabla u, \nabla u^{(\hat{y},\varepsilon)}\rangle\,dx \\ &= -\frac{1}{2}\alpha(1-\kappa)\int_{B_\varepsilon(\hat{y})} \langle \nabla u, \nabla u^{(\hat{y},\varepsilon)}\rangle\,dx\,. \end{aligned} \tag{2.22}$$

Next, we apply Green's formula to $u \in H^1(\Omega)$ and use the fact that u solves the equation $u - f = \alpha \Delta u$ on $B_\varepsilon(\hat{y})$ to obtain that

$$\begin{aligned}
&\alpha \int_{B_\varepsilon(\hat{y})} \langle \nabla u, \nabla u^{(\hat{y},\varepsilon)} \rangle \, dx \\
&\qquad = \alpha \int_{B_\varepsilon(\hat{y})} \langle \nabla (u^{(\hat{y},\varepsilon)} - u), \nabla u \rangle \, dx + \alpha \int_{B_\varepsilon(\hat{y})} |\nabla u|^2 \, dx \\
&\qquad = \int_{B_\varepsilon(\hat{y})} (u^{(\hat{y},\varepsilon)} - u)(f - u) \, dx + \alpha \int_{\partial B_\varepsilon(\hat{y})} (u^{(\hat{y},\varepsilon)} - u) \frac{\partial u}{\partial n} \, ds \\
&\qquad\quad + \alpha \int_{B_\varepsilon(\hat{y})} |\nabla u|^2 \, dx \, .
\end{aligned}$$

To estimate the first integral on the right hand side of above equation, we use the Cauchy-Schwarz inequality and Lemma 2.3 and obtain

$$\begin{aligned}
\Big| \int_{B_\varepsilon(\hat{y})} (u^{(\hat{y},\varepsilon)} - u)(u - f) \, dx \Big| &\le \|u - f\|_{L^2(B_\varepsilon(\hat{y}))} \|u^{(\hat{y},\varepsilon)} - u\|_{L^2(B_\varepsilon(\hat{y}))} \\
&\le 2 \|f\|_{L^\infty(\Omega)} \sqrt{\pi} C_3 \varepsilon^{5/2} \, .
\end{aligned} \tag{2.23}$$

Lemma 2.1 implies that ∇u is Lipschitz continuous on every compact subset of $\Omega \setminus \bar{K}$. Thus there exists some constant c such that

$$\sup_{x \in B_\varepsilon(\hat{y})} \frac{|\nabla u(x) - \nabla u(\hat{y})|}{|x - \hat{y}|} \le c \tag{2.24}$$

for all $\hat{y} \in L$ and $0 < \varepsilon < \operatorname{dist}(L, \Omega \setminus \bar{K})/2$. In particular, we have the estimate

$$\begin{aligned}
&\Big| \int_{B_\varepsilon(\hat{y})} |\nabla u|^2 \, dx - \varepsilon^2 \pi |\nabla u(\hat{y})|^2 \Big| \\
&\qquad\qquad = \Big| \int_{B_\varepsilon(\hat{y})} \langle \nabla u - \nabla u(\hat{y}), \nabla u + \nabla u(\hat{y}) \rangle \, dx \Big| \\
&\qquad\qquad \le c \varepsilon \int_{B_\varepsilon(\hat{y})} |\nabla u + \nabla u(\hat{y})| \, dx \\
&\qquad\qquad \le 2 \, c \, \pi C_1 \|f\|_{H^1(\Omega)} \varepsilon^3 \, .
\end{aligned} \tag{2.25}$$

The change of variable $x = \hat{y} + \varepsilon \tilde{x}$, application of Lemma 2.7 and the property (2.24) yields

$$\begin{aligned}
&\int_{\partial B_\varepsilon(\hat{y})} \big(u^{(\hat{y},\varepsilon)}(x) - u(x) \big) \frac{\partial u}{\partial n}(x) \, ds(x) \\
&\quad = \varepsilon \int_{\partial B(0)} \big(u^{(\hat{y},\varepsilon)}(\hat{y} + \varepsilon \tilde{x}) - u(\hat{y} + \varepsilon \tilde{x}) \big) \frac{\partial u}{\partial n}(\hat{y} + \varepsilon \tilde{x}) \, ds(\tilde{x}) \\
&\quad = \varepsilon^2 \left(\frac{1}{\kappa} - 1 \right) \nabla u(\hat{y})^T \int_{\partial B(0)} \phi(\tilde{x}) \frac{\partial u}{\partial n}(\varepsilon \tilde{x}) \, ds(\tilde{x}) + O(\varepsilon^{5/2}) \\
&\quad = \varepsilon^2 \left(\frac{1}{\kappa} - 1 \right) \nabla u(\hat{y})^T \left(\int_{\partial B(0)} \phi(\tilde{x}) n(\tilde{x})^T ds(\tilde{x}) \right) \nabla u(\hat{y}) + O(\varepsilon^{5/2}) \, ,
\end{aligned} \tag{2.26}$$

where T denotes the vector transpose. Combining the estimates (2.23), (2.25) and (2.26) with (2.21) and (2.22), we obtain

$$\begin{aligned}
&G(u^{(\hat{y},\varepsilon)}, v^{(\hat{y},\varepsilon)}) - G(u,v)\\
&\quad = -\varepsilon^2 \frac{1}{2}\alpha(1-\kappa)\left(\frac{1}{\kappa}-1\right)\nabla u(\hat{y})^T\left(\int_{\partial B}\phi(\tilde{x})n(\tilde{x})^T ds(\tilde{x})\right)\nabla u(\hat{y}) \\
&\qquad - \varepsilon^2\frac{1}{2}\alpha(1-\kappa)\,\pi\,|\nabla u(\hat{y})|^2 + O(\varepsilon^{5/2})\,.
\end{aligned} \tag{2.27}$$

Inserting the explicit formula for the function ϕ, given in (2.20), to the asymptotic expansion (2.27), we get

$$G(u^{(\hat{y},\varepsilon)}, v^{(\hat{y},\varepsilon)}) - G(u,v) = -\varepsilon^2\pi\alpha\frac{1-\kappa}{1+\kappa}\,|\nabla u(\hat{y})|^2 + O(\varepsilon^{5/2})\,,$$

which ends the proof of Theorem 2.1.

3 The Γ-convergence of $J_{\varepsilon,\kappa(\varepsilon)}$ to F

In this section we prove that the sequence of functionals $J_{\varepsilon,\kappa}$ converges to the Mumford–Shah functional in the sense of Γ-convergence, if the parameter κ, depending on ε, tends sufficiently fast to zero as $\varepsilon \to 0$.

Let us first recall the definition of the Γ-limit:

Definition 3.1 *Let X be a topological space and $J_j\colon X \to [0,+\infty]$ a sequence of functionals on X. Denote moreover, for $x \in X$, by $\mathcal{N}(x)$ the set of all open neighborhoods of x. Then the* Γ-lower limit *and the* Γ-upper limit *of J_j are the functionals defined as*

$$\begin{aligned}
(\Gamma\text{-}\liminf_j J_j)(u) &:= \sup_{U\in\mathcal{N}(u)}\ \liminf_j\ \inf_{v\in U} J_j(v)\,,\\
(\Gamma\text{-}\limsup_j J_j)(u) &:= \sup_{U\in\mathcal{N}(u)}\ \limsup_j\ \inf_{v\in U} J_j(v)\,.
\end{aligned}$$

If the Γ-upper and lower limits coincide, we define the Γ-limit by

$$(\Gamma\text{-}\lim_j J_j)(u) = (\Gamma\text{-}\limsup_j J_j)(u) = (\Gamma\text{-}\liminf_j J_j)(u)\,.$$

In metric spaces, the Γ-limit of a sequence of functionals can be characterized by means of the following result:

Lemma 3.1 *Let X be a metric space and $J_j\colon X \to [0,+\infty]$ a sequence of functionals on X. Let moreover $J\colon X \to [0,+\infty]$ and let $\hat{X}$ be a dense subset of $\{u \in X : J(u) < \infty\}$. Assume moreover that for every $u \in X$ there exists a sequence $u_j \to u$ with $u_j \in \hat{X}$ such that $J(u_j) \to J(u)$. Then $J = \Gamma\text{-}\lim_j J_j$, if the following conditions hold:*

1. *For every $u \in X$ and every sequence $u_j \to u$ with $u_j \in X$ we have*

$$J(u) \le \liminf_j J_j(u_j)\,. \tag{3.1}$$

2. *For every $u \in \hat{X}$ and every $\delta > 0$ there exists a sequence $u_j \to u$ with $u_j \in \hat{X}$ for all j such that*

$$J(u) \geq \limsup_j J_j(u_j) - \delta \,. \tag{3.2}$$

Proof. The result follows by combining the standard metric characterization of the Γ-limit (see, e.g., [24, Prop. 8.1]) with a diagonal sequence argument. □

Define

$$F(u,v) := \frac{1}{2}\int_\Omega (u-f)^2\,dx + \frac{\alpha}{2}\int_{\Omega\setminus S_u} |\nabla u|^2\,dx + \beta\mathcal{H}^1(S_u)\,. \tag{3.3}$$

The main result of this section is the following theorem:

Theorem 3.1 *Let F and $J_{\varepsilon,\kappa}$ be as in* (3.3) *and* (1.3)*, respectively. Assume moreover that $\kappa(\varepsilon) = o(\varepsilon)$ as $\varepsilon \to 0$. Then we have for every sequence $\varepsilon_j \to 0$ that*

$$F = \Gamma\text{-}\lim_j J_{\varepsilon_j,\kappa(\varepsilon_j)}\,.$$

We now prove Theorem 3.1 using the methods introduced by Ambrosio and Tortorelli [2], as presented in the notes by Chambolle [15] and the books by Braides [11, 12]. The proof is split into three parts. In the first part, we will prove the lower bound, inequality (3.1), in the one-dimensional case. In the second part, we will extend this result to dimension 2 using the *slicing method* (see, e.g., [11, 12]). In the last part, we will prove the upper bound, inequality (3.2).

3.1 The Lower Bound for $n = 1$

Let $\Omega \subset \mathbb{R}$ be open and bounded, and $f \in L^\infty(\Omega)$. We define the one-dimensional Mumford–Shah functional $\tilde{F}\colon L^2(\Omega)\times L^2(\Omega) \to [0,+\infty]$ as

$$\tilde{F}(u,v) = \begin{cases} \dfrac{1}{2}\displaystyle\int_\Omega (u-f)^2\,dx + \frac{\alpha}{2}\int_{\Omega\setminus S_u} (u')^2\,dx + \beta\mathcal{H}^0(S_u) & \text{if } v \equiv 1\,,\\ +\infty & \text{else.}\end{cases}$$

Because of technicalities of the proof of the two-dimensional case that result from the restriction of the approximating functionals $J_{\varepsilon,\kappa}$ to lines, it is necessary to use a slightly different definition in the one-dimensional case; instead of only covering the edge set with balls of radius ε, we also allow covers with smaller balls. For each finite set $Y = \{y_i : 1 \leq i \leq m\}$ of points in $\mathbb{R}$ we denote by $M_{\varepsilon,\kappa}(Y,\Omega)$ the set of all functions $v\colon \Omega \to \mathbb{R}$ for which there exist a sequence $\{\delta_i\}_{i=1}^m$ of positive numbers smaller than, or equal to, ε, such that for all $x \in \Omega$ we have

$$v(x) = \begin{cases} \kappa & \text{if } x \in \bigcup_{i=1}^m B_{\delta_i}(y_i),\\ 1 & \text{else.}\end{cases}$$

Furthermore we denote by

$$\tilde{m}_{\varepsilon,\kappa}(v,\Omega) := \inf\bigl\{\mathcal{H}^0(Y) : Y \subset \mathbb{R},\ v \in M_{\varepsilon,\kappa}(Y,\Omega)\bigr\}\,. \tag{3.4}$$

As in the two-dimensional case, we define $\tilde{m}_{\varepsilon,\kappa}(v,\Omega) := +\infty$ if $v \notin M_{\varepsilon,\kappa}(Y,\Omega)$ for any $Y \subset \mathbb{R}^2$.

Finally, we define the functional $\tilde{J}_{\varepsilon,\kappa}\colon L^2(\Omega) \times L^2(\Omega) \to [0,+\infty]$ as

$$\tilde{J}_{\varepsilon,\kappa}(u,v) := \frac{1}{2}\int_\Omega (u-f)^2\,dx + \frac{\alpha}{2}\int_\Omega v(u')^2\,dx + \tilde{m}_{\varepsilon,\kappa}(v,\Omega)\ .$$

Let $(u,v) \in L^2(\Omega) \times L^2(\Omega)$, and assume that $\varepsilon_j \to 0$, $u_j \to u$, and $v_j \to v$. We show in the following that

$$\tilde{F}(u,v) \le \liminf_{j\to\infty} \tilde{J}_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j,v_j)\ .$$

We may assume without loss of generality that the sequence $\tilde{J}_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j,v_j)$ converges to some finite number $c < +\infty$; if $c = +\infty$, the claim trivially holds. Moreover we note that the continuity of the norm implies that $\int_\Omega (u_j - f)^2\,dx \to \int_\Omega (u-f)^2\,dx$.

We first define

$$\Sigma := \bigl\{ x \in \Omega : u \notin H^1(x-\delta, x+\delta) \text{ for all } \delta > 0 \bigr\}\ .$$

In the following, we show that Σ is a finite set. Let therefore $s_1 < s_2 < \ldots < s_m \in \Sigma$, and let $\delta > 0$ be such that the sets $(s_i - 2\delta, s_i + 2\delta)$ are disjoint. Because $u \notin H^1(s_i - \delta, s_i + \delta)$ and $u_j \to u$ in $L^2(\Omega)$, the lower semi-continuity of the functional $w \mapsto \int (w')^2\,dx$ implies that

$$\liminf_{j\to\infty} \int_{s_i-\delta}^{s_i+\delta} (u_j')^2\,dx \ge \int_{s_i-\delta}^{s_i+\delta} (u')^2\,dx = +\infty\ .$$

Thus the inequality

$$\begin{aligned} &\limsup_{j\to\infty} \int_{s_i-\delta}^{s_i+\delta} (u_j')^2\,dx \operatorname*{ess\,inf}_{(s_i-\delta,s_i+\delta)} v_j \\ &\qquad\qquad \le \limsup_{j\to\infty} \int_{s_i-\delta}^{s_i+\delta} v_j (u_j')^2\,dx \le \frac{2}{\alpha} \lim_{j\to\infty} \tilde{J}_{\varepsilon_j}(u_j,v_j) = \frac{2c}{\alpha} \end{aligned}$$

implies that

$$\operatorname*{ess\,inf}_{(s_i-\delta,s_i+\delta)} v_j \to 0 \qquad \text{for all } 1 \le i \le m\ . \tag{3.5}$$

Moreover, we have that

$$\limsup_{j\to\infty} \tilde{m}_{\varepsilon_j,\kappa(\varepsilon_j)}(v_j,\Omega) \le \frac{1}{\beta} \lim_{j\to\infty} \tilde{J}_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j,v_j) = \frac{c}{\beta}\ . \tag{3.6}$$

Thus, for j sufficiently large, there exist sets Y_j with $\mathcal{H}^0(Y_j) \le (\beta+1)/c$ such that $v_j \in M_{\varepsilon_j,\kappa(\varepsilon_j)}(Y_j,\Omega)$. In particular, this implies that $v_j \to 1$ with respect to the L^2-norm, showing that $v = 1$. Now the definition of the sets $M_{\varepsilon_j,\kappa(\varepsilon_j)}(Y_j,\Omega)$ and (3.5) imply that for every $1 \le i \le m$ there exists some $x_i^{(j)} \in X_j$ with $B_{\varepsilon_j}(x_i^{(j)}) \cap (s_i - \delta, s_i + \delta) \ne \emptyset$. Because $\varepsilon_j < \delta/2$ for j large enough and the sets $(s_i - 2\delta, s_i + 2\delta)$ are disjoint, it follows that $x_i^{(j)} \ne x_{i'}^{(j)}$ for j large enough and

$i \neq i'$. Consequently, $\mathcal{H}^0(\Sigma) \leq \tilde{m}_{\varepsilon_j,\kappa(\varepsilon_j)}(v_j,\Omega)$ for j large enough. Because by construction $u \in H^1_{\mathrm{loc}}(\Omega \setminus \Sigma)$, it follows that $S_u \subset \Sigma$, and thus

$$\mathcal{H}^0(S_u) \leq \mathcal{H}^0(\Sigma) \leq \liminf_{j\to\infty} \tilde{m}_{\varepsilon_j,\kappa(\varepsilon_j)}(v_j,\Omega) < +\infty \,. \tag{3.7}$$

Next we have to estimate the term $\int_{\Omega\setminus\Sigma_u} (u')^2\,dx$. By assumption, the sequence $(v_j)_{j\in\mathbb{N}}$ converges to v with respect to the L^2-norm. Thus Egoroff's Theorem (see, for instance, [28, 2.3.7]) implies that for every $\gamma > 0$ there exists a set $\Omega_\gamma \subset \Omega \setminus \Sigma$ with $\mathcal{L}^1(\Omega \setminus \Omega_\gamma) \leq \gamma$ and a subsequence v_{j_k} such that v_{j_k} converges uniformly to $v = 1$. Because v_{j_k} takes only the values 1 and $\kappa_{\varepsilon_{j_k}}$ and $\kappa_{\varepsilon_{j_k}} \to 0$, this shows that, in fact, $v_{j_k}(x) = 1$ for $x \in \Omega_\gamma$ and k sufficiently large. Therefore the convergence of u_{j_k} to u implies that

$$\begin{aligned}\int_{\Omega_\gamma} (u')^2\,dx \leq \liminf_{k\to\infty} \int_{\Omega_\gamma} (u'_{j_k})^2\,dx = \liminf_{k\to\infty} \int_{\Omega_\gamma} v_{j_k}(u'_{j_k})^2\,dx \\ \leq \frac{2}{\alpha} \lim_{k\to\infty} \tilde{J}_{\varepsilon_{j_k}}(u_{j_k},v_{j_k}) = \frac{2c}{\alpha}\,.\end{aligned}$$

Taking the supremum with respect to γ, it follows that $u \in H^1(\Omega \setminus S_u)$ and

$$\int_{\Omega\setminus S_u} (u')^2\,dx \leq \limsup_{j\to\infty} \int_\Omega v_j(u'_j)^2\,dx\,,$$

which concludes the proof.

3.2 The Lower Bound for $n = 2$

The second part of the proof of Theorem 3.1 is concerned with showing (3.1) for $\Omega \subset \mathbb{R}^2$. The proof applies the slicing method following Braides [11, 12]. To that end it is necessary to introduce some notational preliminaries:

We denote for every direction $\xi \in \mathbb{S}^1 := \{x \in \mathbb{R}^2 : |x| = 1\}$ by $\Pi_\xi := \{y \in \mathbb{R}^2 : \langle y,\xi\rangle = 0\}$ the hyperplane passing through 0 that is orthogonal to ξ. If $A \subset \Omega$ is open, we denote by $A_{\xi,y} := \{t \in \mathbb{R} : y + t\xi \in A\} \subset \mathbb{R}$ the one-dimensional slice of A indexed by $y \in \Pi_\xi$. Finally, for all w defined on Ω, we define the one-dimensional function $w^{\xi,y}(t) = w(y + t\xi)$ as the restriction of w to $\Omega_{\xi,y}$.

Next we define for every open set $A \subset \Omega$ a *localized* functional $J_{\varepsilon,\kappa}(u,v,A)$. To that end, we first localize the functional $m_{\varepsilon,\kappa}$. We define

$$m_{\varepsilon,\kappa}(v,A) := \inf\bigl\{\mathcal{H}^0(Y) : Y \subset \mathbb{R}^2,\ v|_A = v_{Y,\kappa}|_A\bigr\}\,.$$

Then we define

$$J_{\varepsilon,\kappa}(u,v,A) := \frac{1}{2}\int_A (u-f)^2 + \alpha v|\nabla u|^2\,dx + 2\varepsilon\beta m_{\varepsilon,\kappa}(v,A)\,.$$

Moreover, we define for each $\xi \in \mathbb{S}^1$ the directional functional

$$J^\xi_{\varepsilon,\kappa}(u,v,A) := \frac{1}{2}\int_A (u-f)^2 + \alpha v\langle\xi,\nabla u\rangle^2\,dx + 2\varepsilon\beta m_{\varepsilon,\kappa}(v,A)\,.$$

Finally, we consider for each $\xi \in \mathbb{S}^1$, $y \in \Pi_\xi$, and $I \subset \Omega_{\xi,y}$ open the one-dimensional functionals

$$F^{\xi,y}(\hat{u}, I) = \frac{1}{2} \int_{I \setminus S_{\hat{u}}} (\hat{u} - f^{\xi,y})^2 + (\hat{u}')^2 \, dx + \beta \mathcal{H}^0(S_{\hat{u}})$$

and

$$J^{\xi,y}_{\varepsilon,\kappa}(\hat{u}, \hat{v}, I) = \frac{1}{2} \int_I (\hat{u} - f^{\xi,y})^2 + \alpha \hat{v} (\hat{u}')^2 \, dx + \beta \tilde{m}_{\varepsilon,\kappa}(\hat{v}, I) \,,$$

where $\tilde{m}_{\varepsilon,\kappa}$ is as defined in (3.4).

We claim that for every u, $v \in L^2(\Omega)$, $\varepsilon > 0$, $0 < \kappa < 1$, $A \subset \Omega$ open, and $\xi \in \mathbb{S}^1$ the inequalities

$$J_{\varepsilon,\kappa}(u, v, A) \geq J^{\xi}_{\varepsilon,\kappa}(u, v, A) \geq \int_{\Pi_\xi} J^{\xi,y}_{\varepsilon,\kappa}(u^{\xi,y}, v^{\xi,y}, A_{\xi,y}) \, d\mathcal{H}^1(y) \tag{3.8}$$

hold. Indeed, the first inequality is a direct consequence of the definition of the involved functionals. For the second inequality, note first that, by Fubini's theorem,

$$\int_A (u - f)^2 + \alpha v \langle \xi, \nabla u \rangle^2 \, dx = \int_{\Pi^\xi} \int_{A_{\xi,y}} (u^{\xi,y} - f^{\xi,y})^2 + \alpha v^{\xi,y} (u^{\xi,y})'^2 \, dt \, d\mathcal{H}^1(y) \,.$$

Thus it remains to show that

$$2\varepsilon \, m_{\varepsilon,\kappa}(v, A) \geq \int_{\Pi_\xi} \tilde{m}_{\varepsilon,\kappa}(v^{\xi,y}, A_{\xi,y}) \, d\mathcal{H}^1(y) \tag{3.9}$$

whenever $v \in L^2(A)$. In case $m_{\varepsilon,\kappa}(v, A) = +\infty$, this inequality trivially holds. Else, there exists a set $Y = \{y_1, \ldots, y_m\} \subset \mathbb{R}^2$ with $m = m_{\varepsilon,\kappa}(v, A)$ such that

$$v(x) = \begin{cases} \kappa_\varepsilon & \text{if } x \in \bigcup_{i=1}^m B_\varepsilon(y_i) \,, \\ 1 & \text{else.} \end{cases}$$

Then

$$\begin{aligned} 2\varepsilon \, m &= \sum_{i=1}^m \mathcal{H}^1\bigl(\{y \in \Pi_\xi : B_\varepsilon(y_i) \cap (y + \mathbb{R}\xi) \neq \emptyset\}\bigr) \\ &\geq \sum_{i=1}^m \mathcal{H}^1\bigl(\{y \in \Pi_\xi : B_\varepsilon(y_i) \cap (y + \mathbb{R}\xi) \cap A \neq \emptyset\}\bigr) \\ &= \int_{\Pi_\xi} \mathcal{H}^0\bigl(\{i : B_\varepsilon(y_i) \cap (y + \mathbb{R}\xi) \cap A \neq \emptyset\}\bigr) \, d\mathcal{H}^1(y) \,. \end{aligned}$$

Moreover the definition of $\tilde{m}_{\varepsilon,\kappa}(v^{\xi,y}, A_{\xi,y})$ implies that

$$\mathcal{H}^0\bigl(\{i : B_\varepsilon(y_i) \cap A_{\xi,y} \neq \emptyset\}\bigr) \geq \tilde{m}_{\varepsilon,\kappa}(v^{\xi,y}, A_{\xi,y})$$

for all y and ξ. This shows (3.9), which in turn implies (3.8).

Now let $(u, v) \in L^2(\Omega) \times L^2(\Omega)$, and assume that $\varepsilon_j \to 0$, $u_j \to u$, and $v_j \to v$. As in the one-dimensional case we have to show that

$$F(u, v) \leq \liminf_{j \to \infty} J_{\varepsilon_j, \kappa(\varepsilon_j)}(u_j, v_j) \,.$$

Again, we assume without loss of generality that the sequence $J_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j,v_j)$ converges to some finite number $c < +\infty$; the claim being trivial if $c = +\infty$. In particular, this implies that $v = 1$ almost everywhere. Now (3.8), Fatou's Lemma, and the Γ-convergence result for the one-dimensional case imply that, for each open set $A \subset \Omega$ and each direction $\xi \in \mathbb{S}^1$, we have

$$\begin{aligned}
&\liminf_{j\to\infty} J_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j,v_j,A)\\
&\qquad\geq \liminf_{j\to\infty} \int_{\Pi_\xi} J^{\xi,y}_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j^{\xi,y}, v_j^{\xi,y}, A_{\xi,y})\,d\mathcal{H}^1(y)\\
&\qquad\geq \int_{\Pi_\xi} \liminf_{j\to\infty} J^{\xi,y}_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j^{\xi,y}, v_j^{\xi,y}, A_{\xi,y})\,d\mathcal{H}^1(y)\\
&\qquad\geq \int_{\Pi_\xi} F^{\xi,y}(u^{\xi,y}, A_{\xi,y})\,d\mathcal{H}^1(y)\\
&\qquad= \frac{1}{2}\int_{A\setminus S_u} (u-f)^2 + \alpha\langle\xi,\nabla u\rangle^2\,dx\\
&\qquad\qquad + \beta\int_{\Pi_\xi} \mathcal{H}^0(S_{u^{\xi,y}} \cap A_{\xi,y})\,d\mathcal{H}^1(y)\\
&\qquad= \frac{1}{2}\int_{A\setminus S_u} (u-f)^2 + \alpha\langle\xi,\nabla u\rangle^2\,dx + \beta\int_{S_u\cap A} |\langle\xi,\nu_u\rangle|\,d\mathcal{H}^1(x)\ .
\end{aligned}$$

Now let $(\xi_i)_{i\in\mathbb{N}} \subset \mathbb{S}^1$ be a dense subset. Then [11, p. 191] implies that

$$\begin{aligned}
\liminf_{j\to\infty} J_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j,v_j) &\geq \frac{1}{2}\int_{\Omega\setminus S_u} (f-u)^2 + \alpha\sup_i\langle\xi_i,\nabla u\rangle^2\,dx\\
&\qquad + \int_{S_u} \sup_i|\langle\xi_i,\nu_u\rangle|\,d\mathcal{H}^{n-1}(y)\\
&= F(u)\ .
\end{aligned}$$

3.3 The Upper Bound

We now prove inequality (3.2). To that end, we consider the set $\mathcal{W}(\Omega)$ consisting of all functions $u \in SBV(\Omega)$ for which the following hold:

1. $\mathcal{H}^1(\overline{S}_u \setminus S_u) = 0$.
2. The set $\overline{S}_u$ is the union of a finite number of almost disjoint line segments contained in Ω.
3. $u|_{\Omega\setminus\bar{S}_u} \in W^{1,\infty}(\Omega \setminus S_u)$.

Obviously, this set is dense in $SBV(\Omega)$ with respect to the L^2-topology. Moreover, it has been shown in [22, 23] that, for every $u \in SBV(\Omega)$, there exists a sequence $u_j \to u$ with $u_j \in \mathcal{W}(\Omega)$ for every j such that $F(u_j) \to F(u)$. Using Lemma 3.1, for proving (3.2), we therefore have to find for every $u \in \mathcal{W}(\Omega)$, $\delta > 0$, and $\varepsilon_j \to 0$ sequences $u_j \to 0$, $v_j \to 1$ as $j \to \infty$, such that $\limsup_{j\to\infty} J_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j,v_j) \leq F(u) + \delta$.

Let therefore $u \in \mathcal{W}(\Omega)$ be fixed. By definition of $\mathcal{W}(\Omega)$, there exist $k \in \mathbb{N}$ and a_i, $b_i \in \Omega$, $1 \le i \le k$, such that $\overline{S}_u = \bigcup_{i=1}^k [a_i, b_i]$. Moreover,

$$\mathcal{H}^1(S_u) = \mathcal{H}^1(\overline{S}_u) = \sum_{i=1}^k \|b_i - a_i\| \, .$$

Now define for $\varepsilon > 0$ and $0 < c < 1$

$$K(\varepsilon, c) := \bigl\{ x \in \Omega : \operatorname{dist}(x, S_u) < c\varepsilon \bigr\} \, .$$

Let moreover $\mu(\varepsilon, c) \in \mathbb{N}$ and $x_l^{(\varepsilon,c)} \in \Omega$, $1 \le l \le \mu(\varepsilon, c)$, be such that

$$K(\varepsilon, c) \subset \bigcup_{l=1}^{\mu(\varepsilon,c)} B_\varepsilon(x_l^{(\varepsilon,c)}) \, .$$

Now note that, if we place the centers of the balls on a line segment $[a_i, b_i]$, then they cover the whole set $\bigl\{ x \in \Omega : \operatorname{dist}(x, [a_i, b_i]) < c\varepsilon \bigr\}$ provided that the distance between two adjacent centers is at most $2\varepsilon\sqrt{1-c^2}$. Thus it follows that one can cover each set $\bigl\{ x \in \Omega : \operatorname{dist}(x, [a_i, b_i]) < c\varepsilon \bigr\}$ with at most $\frac{\|b_i - a_i\|}{2\varepsilon\sqrt{1-c^2}} + 1$ balls of radius ε. Consequently, we can choose the centers x_l in such a way that

$$\mu(\varepsilon, c) \le k + \sum_{i=1}^k \frac{\|b_i - a_i\|}{2\varepsilon\sqrt{1-c^2}} = k + \frac{\mathcal{H}^1(S_u)}{2\varepsilon\sqrt{1-c^2}} \, .$$

Let now

$$v^{(\varepsilon,c)}(x) := \begin{cases} \kappa(\varepsilon) & \text{if } x \in \bigcup_{l=1}^{\mu(\varepsilon,c)} B_\varepsilon(x_l^{(\varepsilon,c)}) \, , \\ 1 & \text{else.} \end{cases}$$

Then, for every c we have $v^{(\varepsilon,c)} \to 1$ as $\varepsilon \to 0$. Moreover,

$$m_{\varepsilon,\kappa(\varepsilon)}(v^{(\varepsilon,c)}, \Omega) \le \mu(\varepsilon, c) \, .$$

Define now

$$u^{(\varepsilon,c)}(x) := u(x) \, \min \left(\frac{\operatorname{dist}(x, S_u)}{c\varepsilon}, 1 \right) \, .$$

Then $u^{(\varepsilon,c)}(x) = u(x)$ for $x \notin K(\varepsilon, c)$ and $u^{(\varepsilon,c)} \to u$ as $\varepsilon \to 0$. Denoting $d(x) = \operatorname{dist}(x, S_u)$, we have for almost every $x \in K(\varepsilon, c)$

$$\nabla u^{(\varepsilon,c)}(x) = \nabla u(x) \frac{d(x)}{c\varepsilon} + u(x) \frac{\nabla d(x)}{c\varepsilon} \, .$$

Thus the triangle inequality and the fact that $|\nabla d(x)| = 1$ almost everywhere imply that

$$|\nabla u^{(\varepsilon,c)}(x)| \le |\nabla u(x)| \frac{d(x)}{c\varepsilon} + |u(x)| \frac{|\nabla d(x)|}{c\varepsilon} \le |\nabla u(x)| + \frac{\|u\|_\infty}{c\varepsilon}$$

for almost every $x \in K(\varepsilon, c)$. Therefore, for almost every $x \in K(\varepsilon, c)$,

$$|\nabla u^{(\varepsilon,c)}(x)|^2 \le 2\|\nabla u\|_\infty^2 + 2\frac{\|u\|_\infty^2}{c^2\varepsilon^2} \, . \tag{3.10}$$

Now consider each term of the functional $J_{\varepsilon,\kappa(\varepsilon)}$ separately. We have

$$\int_\Omega (u^{(\varepsilon,c)} - f)^2 dx = \int_{\Omega\setminus K(\varepsilon,c)} (u-f)^2 dx + \int_{K(\varepsilon,c)} (u^{(\varepsilon,c)} - f)^2 dx$$

$$\to_{\varepsilon\to 0} \int_\Omega (u-f)^2 dx \,.$$

From (3.10) we get

$$\int_\Omega v^{(\varepsilon,c)} |\nabla u^{(\varepsilon,c)}|^2 dx \le \int_{\Omega\setminus K(\varepsilon,c)} |\nabla u|^2 dx + \int_{K(\varepsilon,c)} \kappa(\varepsilon) |\nabla u^{(\varepsilon,c)}|^2 dx$$

$$\le \int_{\Omega\setminus K(\varepsilon,c)} |\nabla u|^2 dx + 2\kappa(\varepsilon)\|\nabla u\|_\infty^2 \mathcal{L}^2(K(\varepsilon,c)) + \frac{2\kappa(\varepsilon)\|u\|_\infty^2 \mathcal{L}^2(K(\varepsilon,c))}{c^2\varepsilon^2} \,.$$

Because $\kappa(\varepsilon) = o(\varepsilon)$ as $\varepsilon \to 0$ and

$$\mathcal{L}^2(K(\varepsilon,c)) \le \mathcal{L}^2(K(\varepsilon,1)) \le 2\varepsilon\mathcal{H}^1(S_u) + k\pi\varepsilon^2 = O(\varepsilon) \qquad \text{as } \varepsilon \to 0\,,$$

it follows that

$$\limsup_{\varepsilon\to 0} \int_\Omega v^{(\varepsilon,c)} |\nabla u^{(\varepsilon,c)}|^2 dx \le \int_{\Omega\setminus S_u} |\nabla u|^2 \, dx \,.$$

Finally, the construction of $v^{(\varepsilon,c)}$ implies that

$$2\varepsilon\, m_{\varepsilon,\kappa(\varepsilon)}(v^{(\varepsilon,c)}, \Omega) \le 2k\varepsilon + \frac{\mathcal{H}^1(S_u)}{\sqrt{1-c^2}} \,.$$

Let now $\varepsilon_j \to 0$ as $j \to \infty$ and define $u_j := u^{(\varepsilon_j,c)}$, $v_j := v^{(\varepsilon_j,c)}$. Then it follows that

$$\limsup_{j\to\infty} J_{\varepsilon_j,\kappa(\varepsilon_j)}(u_j, v_j) \le \frac{1}{2}\int_\Omega (u-f)^2 \, dx + \frac{\alpha}{2}\int_{\Omega\setminus S_u} |\nabla u|^2 \, dx + \frac{\beta}{\sqrt{1-c^2}}\mathcal{H}^1(S_u) \,.$$

Since $0 < c < 1$ was arbitrary and $\mathcal{H}^1(S_u) < \infty$, we obtain (3.2) with $\delta = (1 - 1/\sqrt{1-c^2})\mathcal{H}^1(S_u)$, which concludes the proof of Theorem 3.1.

4 Numerical Implementation

4.1 Proposed Algorithm

Based on Theorem 2.1, we propose the following algorithm for the approximate minimization of the functional $J_{\varepsilon,\kappa}$ for fixed $\varepsilon > 0$ and $\kappa > 0$.

Algorithm 4.1 *Let $f \in L^\infty(\Omega)$, α, $\beta > 0$, $\varepsilon > 0$, and $0 < \kappa < 1$ be given.*

Set $k = 0$ and $K_0 := \emptyset$.

1. *Define*

$$v_k(x) := \begin{cases} \kappa & \text{if } x \in K_k\,, \\ 1 & \text{if } x \in \Omega \setminus K_k\,. \end{cases}$$

2. *Define*
$$u_k := \arg\min_u G(u, v_k)\ .$$
3. *Find* $y \in \Omega \setminus K_k$ *such that* $|\nabla u_k(y)|$ *is maximal.*
4. *If*
$$\frac{\alpha}{2}\,\pi\,\frac{1-\kappa}{1+\kappa}|\nabla u_k(y)|^2 < \frac{\beta}{\varepsilon}\,, \tag{4.1}$$
stop.

   Else set $K_{k+1} := K_k \cup B_\varepsilon(y)$*, increase* k *by one and go to step 1.*

Steps 3 and 4 of the algorithm use the results of Theorem 2.1. This theorem states that adding in the k-th step a point y to the edge indicator will approximately result in a decrease of the functional G by approximately $\varepsilon^2\alpha\pi(1+\kappa)|\nabla u_k(y)|^2/(1-\kappa)$. Thus, we will obtain the steepest descent, if we add a point y, where $|\nabla u_k(y)|$ is maximal. At the same time the adding of another ball leads to an increase of the term $m_{\varepsilon,\kappa}$ by $2\beta\varepsilon$. In total, the value of $J_{\varepsilon,\kappa}$ will increase if (4.1) holds, else $J_{\varepsilon,\kappa}$ will decrease and therefore it makes sense to include the point y into the edge set.

Remark 4.1 *In order to increase the performance of the algorithm, it makes sense to add not just one ball in each iteration, but rather several ones. Also in this case a similar approximation as Theorem 2.1 holds, and thus the same criterion for adding new points can be applied. This strategy has been used in the numerical examples below.*

4.2 Numerical Results

We now compare the results obtained with Algorithm 4.1 with results obtained using the approximation introduced by Ambrosio and Tortorelli [3]. This latter method consists in minimizing the functional

$$I_\varepsilon(u,v) := \frac{1}{2}\int_\Omega (u-f)^2\,dx + \frac{\alpha}{2}\int_\Omega v^2|\nabla u|^2\,dx + \frac{1}{2}\int_\Omega \Big(\varepsilon|\nabla u|^2 + \frac{1}{4\varepsilon}(v-1)^2\Big)\,dx\ . \tag{4.2}$$

Again, the function v serves as an edge indicator in the sense that the points where v is close to zero are an approximation of the edge set K of the solution of the Mumford–Shah functional. In contrast to the approximation by means of the functional $J_{\varepsilon,\kappa}$, however, where the edge set is given as the points where the function v is equal to κ, in case of the functional (4.2) one has to threshold v in order to obtain a precisely defined edge set.

The minimization of I_ε has been carried out by alternately solving the corresponding Euler–Lagrange equations with respect to u and v. For the discretization, we have used a finite element approach with piecewise bilinear basis functions on the pixel grid. The same discretization has been used for the computation of u_k in the second step of Algorithm 4.1.

Figures 1 and 2 show a comparison of the results of the Ambrosio–Tortorelli approximation and Algorithm 4.1. The edge indicators are in both cases comparable, though our algorithm in general classifies more points as edges. The main difference between the results is that the Ambrosio–Tortorelli approximation leads to a diffuse edge indicator, while our method produces well defined edges. As a consequence, also the smoothed images tend to be less blurred; compare, for instance, the various light reflections in Figure 2.

Acknowledgement

The work of OS has been supported by the Austrian Science Fund (FWF) within the national research networks Industrial Geometry, project 9203-N12, and Photoacoustic Imaging in Biology and Medicine, project S10505-N20. The authors thank Prof. Helmut Neunzert for his continuous encouragement of this collaboration.

References

[1] R. A. Adams. *Sobolev Spaces.* Academic Press, New York, 1975.

[2] L. Ambrosio. Existence theory for a new class of variational problems. *Arch. Ration. Mech. Anal.*, 111:291–322, 1990.

[3] L. Ambrosio and V. M. Tortorelli. Approximation of functionals depending on jumps by elliptic functionals via Γ-convergence. *Comm. Pure Appl. Math.*, 43(8):999–1036, 1990.

[4] H. Ammari and H. Kang *Reconstruction of small inhomogeneities from boundary measurements*, volume 1846 of *Lecture Notes in Mathematics.* Springer, 2004.

[5] S. Amstutz. Sensitivity analysis with respect to a local perturbation of the material property. *Asymptotic Anal.*, 49(1-2):87–108, 2006.

[6] G. Aubert, L. Blanc-Féraud, and R. March. An approximation of the Mumford-Shah energy by a family of discrete edge-preserving functionals. *Nonlinear Anal.*, 64(9):1908–1930, 2006.

[7] G. Aubert and P. Kornprobst. *Mathematical Problems in Image Processing: Partial Differential Equations and the Calculus of Variations (second edition)*, volume 147 of *Applied Mathematical Sciences.* Springer-Verlag, 2006.

[8] D. Auroux, L. J. Belaid, and M. Masmoudi. A topological asymptotic analysis for the regularized grey-level image classification problem.
Math. Model. Numer. Anal., 41(3), 2007.

[9] D. Auroux and M. Masmoudi. Image processing by topological asymptotic expansion. *J. Math. Imaging Vision*, 33(2), 2009.

[10] A. Blake and A. Zisserman. *Visual Reconstruction*, MIT Press, 1987.

[11] A. Braides. *Approximation of free-discontinuity problems*, volume 1694 of *Lecture Notes in Mathematics.* Springer-Verlag, Berlin, 1998.



Fig. 1: *Upper left:* Original image. *Upper right:* Smoothing with the Ambrosio–Tortorelli Approximation. *Middle left:* Smoothing with Algorithm 4.1. *Middle right:* Edge indicator for the Ambrosio–Tortorelli Approximation. *Lower left:* Edge indicator for Algorithm 4.1. *Lower right:* Edge indicator for the Ambrosio–Tortorelli Approximation, thresholded at 0.8. For both methods, $\alpha = 20$, $\beta = 200$, $\varepsilon = 0.05$.



Fig. 2: *Upper left:* Original image. *Upper right:* Smoothing with the Ambrosio–Tortorelli Approximation. *Middle left:* Smoothing with Algorithm 4.1. *Middle right:* Edge indicator for the Ambrosio–Tortorelli Approximation. *Lower left:* Edge indicator for Algorithm 4.1. *Lower right:* Edge indicator for the Ambrosio–Tortorelli Approximation, thresholded at 0.9. For both methods, $\alpha = 1$, $\beta = 500$, $\varepsilon = 0.05$.

[12] A. Braides. Γ-*convergence for beginners*, volume 22 of *Oxford Lecture Series in Mathematics and its Applications.* Oxford University Press, Oxford, 2002.

[13] A. Braides and G. Dal Maso. Non-local approximation of the Mumford-Shah functional. *Calc. Var. Partial Differential Equations*, 5:293–322, 1997.

[14] D.J. Cedio-Fengya, S. Moskow and M.S. Vogelius. Identification of conductivity imperfections of small diameter by boundary measurements. Continuous dependence and computational reconstruction. *Inverse Problems*, 14(3):553–595, 1998.

[15] A. Chambolle. *Mathematical problems in image processing.* ICTP Lecture Notes, II. Abdus Salam International Centre for Theoretical Physics, Trieste, 2000. Inverse problems in image processing and image segmentation: some mathematical and numerical aspects, Available electronically at `http://www.ictp.trieste.it/~pub_off/lectures/vol2.html`.

[16] A. Chambolle. Image segmentation by variational methods: Mumford-Shah functional and the discrete approximations. *SIAM J. Appl. Math.*, 55(3):827–863, 1995.

[17] A. Chambolle. Finite-differences discretizations of the Mumford-Shah functional. *M2AN Math. Model. Numer. Anal.*, 33(2):261–288, 1999.

[18] A. Chambolle and G. Dal Maso. Discrete approximation of the Mumford-Shah functional in dimension two. *M2AN Math. Model. Numer. Anal.*, 33(4):651–672, 1999.

[19] T. F. Chan and L. A. Vese. Active contours without edges. *IEEE Trans. Image Processing*, 10(2):266–277, 2001.

[20] D. Colton and R. Kress. *Integral Equation Methods in Scattering Theory.* Wiley, New York, 1983.

[21] D. Colton and R. Kress. *Inverse Acoustic and Electromagnetic Scattering Theory.* Springer, 1998.

[22] G. Cortesani. Strong approximation of GSBV functions by piecewise smooth functions. *Ann. Univ. Ferrara Sez. VII (N.S.)*, 43:27–49 (1998), 1997.

[23] G. Cortesani and R. Toader. A density result in SBV with respect to non-isotropic energies. *Nonlinear Anal.*, 38(5, Ser. B: Real World Appl.):585–604, 1999.

[24] G. Dal Maso. *An Introduction to* Γ-*Convergence*, volume 8 of *Progress in Nonlinear Differential Equations and their Applications.* Birkhäuser, 1993.

[25] G. Dal Maso, J. M. Morel, and S. Solimini. A variational method in image segmentation: Existence and approximation results. *Acta Mathematica*, 168(1):89–151, 1992.

[26] E. De Giorgi, M. Carriero, and A. Leaci. Existence theorem for a minimum problem with free discontinuity set. *Arch. Ration. Mech. Anal.*, 108:195–218, 1989.

[27] H. W. Engl, M. Hanke, and A. Neubauer. *Regularization of inverse problems*, volume 375 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1996.

[28] H. Federer. *Geometric Measure Theory*. Classics in Mathematics. Springer Verlag, Berlin, 1996. Repr. of the 1969 ed.

[29] R. A. Feijóo, A. Novotny, C. Padra, and E. Taroco. The topological derivative for the Poisson problem. *Math. Mod. Meth. Appl. Sci.*, 13(12):1825–1844, 2003.

[30] S. Garreau, P. Guillaume, and M. Masmoudi. The topological asymptotic for PDE systems: The elasticity case. *SIAM J. Control Optimiz*, 39(6):1756–1778, 2000.

[31] D. Gilbarg and N. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Classics in Mathematics. Springer Verlag, Berlin, 2001. Reprint of the 1998 edition.

[32] G. Koepfler, C. Lopez, and J. M. Morel. A multiscale algorithm for image segmentation by variational method. *SIAM J. Numer. Anal.*, 31(1):282–299, 1994.

[33] S. G. Krein and Yu. I. Petunin. Scales of Banach spaces. *Russ. Math. Surv.*, 21(85):85–159, 1966.

[34] R. Kress. *Linear Integral Equations*. Springer, 1989.

[35] J.-M. Morel and S. Solimini. *Variational Methods in Image Segmentation*. Birkhäuser, Boston, 1995.

[36] D. Mumford and J. Shah. Optimal approximations by piecewise smooth functions and associated variational problems. *Comm. Pure Appl. Math.*, 42(5):577–685, 1989.

[37] M. Muszkieta. Optimal edge detection by topological asymptotic analysis. *Math. Mod. Meth. Appl. Sci.*, 19(11):2127–2143, 2009.

[38] P. D. Polyanin. *Handbook of Linear Partial Differential Equations*. Chapman & Hall, 2002.

[39] O. Scherzer, editor. *Handbook of Mathematical Methods in Imaging*. Springer-Verlag New York Inc., 2010.

[40] O. Scherzer, M. Grasmair, H. Grossauer, M. Haltmeier, and F. Lenzen. *Variational methods in imaging*, volume 167 of *Applied Mathematical Sciences*. Springer, New York, 2009.

[41] J. Sokołowski and A. Żochowski. On topological derivative in shape optimization. *SIAM J. Control Optimiz*, 37(4):1251–1272, 1999.

[42] J. Spanier. *An Atlas of Functions.* Taylor & Francis, 1987.

[43] M. S. Vogelius and D. Volkov. Asymptotic formulas for perturbations in the electromagnetic fields due to the presence of inhomogeneities of small diameter. *Math. Model. Numer. Anal.*, 34(4):723–748, 2000.